# Enhancing the assimilation of microwave water vapor channels

# Isaac Moradi<sup>1</sup> and Ralph Ferraro<sup>2</sup>

1. Earth System Science Interdisciplinary Center (ESSIC),
University of Maryland, College Park, MD

2. SCSB, STAR, NESDIS, NOAA, College Park, MD

NASA AIRS Science Team Meeting Caltech, Pasadena, CA, USA March 22-24, 2016

March 22, 2016



## Outline

- Variational Bias Correction
- Radiometric Error
- Validation
- Assimilation of All-Weather MW
- Conclusions

# Variational Bias Correction

### Variational Bias Correction

Cost function for 3D-Var Data Assimilation:

$$J(\vec{x}) = \overbrace{\frac{1}{2}(\vec{x} - \vec{x_b})^T \vec{B}^{-1}(\vec{x} - \vec{x_b})}^{J_b} + \overbrace{\frac{1}{2}(H(\vec{x}) - \vec{y})^T \vec{R}^{-1}(H(\vec{x}) - \vec{y})}^{J_o}$$

$$y = Tb + \epsilon_r + \epsilon_s$$


 $\epsilon_s$  is the random error (R) and  $\epsilon_s$  is known as observation bias or representativeness error that is taken into account using the variational bias correction:

$$\epsilon_s = \sum\limits_{k=1}^{N} eta_k p_k + b^{angle}$$


The control variables  $(p_k)$  include cloud liquid water (CLW); temperature lapse rate; and the square of the temperature lapse rate.




### Variational Bias Correction




# NOAA GDAS Operational System






# **NASA GMAO**




## **ECMWF**

### All MHS MetOp-A Observations




# Assimilated MHS MetOp-A



### Issues in Variational Bias Correction?


$$\epsilon_s = Obs_e + RT_e + Col_e$$

- variational bias correction does not distinguish between the model error and the observations error
- in variational bias correction, the RT and collocation errors are counted as observations error
- biased NWP fields or some biased observations could lead to the over-correction of good observations




### Collocation Error


# Collocation error as a function of distance



# Collocation Error vs. the time difference




# Clouds and Surface Emissivity




# Radiometric Error

# Quantifying Radiometric Error


# Collocated observations from similar channels




# Time series of the difference between the observations



# Level 1b MHS/AMSU-B




# Corrected MHS/AMSU-B




# Validation


# L1b Bias Correction




# L1b Bias Correction




# L1b Bias Correction




# GMAO MetOp-A MHS Tb's



### **GMAO Total Bias Correction**




### **GMAO Total Bias Correction**



# Assimilation of All-Weather MW

# Assimilation of All-Weather MW Using BMCI



Black: observed, Red: bias-corrected FG for AMSR-E 37v. C37 is the cloud index derived from polarization difference (one minus v-h all sky divided by v-h clear) C37 varies from (for clear-sky to 1 for opaque clouds (Geer and Bauer, 2010).

# Conclusions

### Conclusions

- variational bias correction technique does not distinguish between error sources - errors may compensate for each other
- variational bias correction does not especially work for water vapor channels because of large error in the NWP water vapor fields
- more robust and physical bias correction techniques are available that can quantify different components of the representativeness error
- some preliminary results are presented but more work is required to properly validate the impact of bias corrected observations on the system

# Thank you for your attention!